

**Drug-Resistant Tuberculosis Cases,
North Carolina 2014**

County	INH Resistance	INH & RIF Resistance
Alamance	0	0
Alexander	0	0
Alleghany	0	0
Anson	0	0
Ashe	0	0
Avery	0	0
Beaufort	0	0
Bertie	0	0
Bladen	0	0
Brunswick	0	0
Buncombe	0	0
Burke	0	0
Cabarrus	0	0
Caldwell	0	0
Camden	0	0
Carteret	1	0
Caswell	0	0
Catawba	0	0
Chatham	0	0
Cherokee	0	0
Chowan	0	0
Clay	0	0
Cleveland	0	0
Columbus	0	0
Craven	1	0
Cumberland	0	0
Currituck	0	0
Dare	0	0
Davidson	0	0
Davie	0	0
Duplin	0	0
Durham	1	0
Edgecombe	0	0
Forsyth	0	0
Franklin	0	0
Gaston	0	0
Gates	0	0
Graham	0	0
Granville	0	0
Greene	0	0
Guilford	0	1
Halifax	0	0
Harnett	0	0
Haywood	0	0
Henderson	1	0
Herford	0	0
Hoke	0	0
Hyde	0	0
Iredell	0	0
Jackson	0	0
Johnston	0	0
Jones	0	0
Lee	0	0
Lenoir	1	0
Lincoln	0	0
Macon	0	0

Madison	0	0
Martin	0	0
McDowell	0	0
Mecklenburg	2	0
Mitchell	0	0
Montgomery	0	0
Moore	0	0
Nash	0	0
New Hanover	0	0
Northampton	0	0
Onslow	0	0
Orange	0	0
Pamlico	0	0
Pasquotank	0	0
Pender	0	0
Perquimans	0	0
Person	0	0
Pitt	1	0
Polk	0	0
Randolph	0	0
Richmond	0	0
Robeson	0	0
Rockingham	0	0
Rowan	0	0
Rutherford	0	0
Sampson	0	0
Scotland	0	0
Stanly	0	0
Stokes	0	0
Surry	0	0
Swain	0	0
Transylvania	0	0
Tyrrell	0	0
Union	0	0
Vance	0	0
Wake	0	0
Warren	0	0
Washington	0	0
Watagua	0	0
Wayne	1	0
Wilkes	0	0
Wilson	0	0
Yadkin	0	0
Yancey	0	0
North Carolina	9	1

INH resistance: The TB bacteria are resistant to isoniazid (INH) rendering INH no longer effective for treatment. INH resistance may include resistance to additional drugs, but excludes resistance to Rifampin (RIF). Resistance may be found at initial or final testing.

Source: NC TB Control Program

Date: Feb 13, 2015