



NC DEPARTMENT OF
**HEALTH AND
HUMAN SERVICES**

ROY COOPER • Governor
KODY H. KINSLEY • Secretary
Mark T. Benton • Deputy Secretary for Health
Kelly Kimple • Acting Director, Division of Public Health

To: All North Carolina Clinicians
From: Emma Doran, MD, MPH, Medical Epidemiologist
Subject: 2024-2025 Respiratory Virus Season: **Testing Update for NC Clinicians** (3 pages)
Date: September 25, 2024

This memo provides information and guidance to NC clinicians regarding testing for respiratory viruses in North Carolina during the 2024-2025 respiratory season. As guidance may change during the season, up to date information will be available at flu.nc.gov. Testing plays a critical role in detecting acute respiratory viral infections including infections with novel or variant influenza viruses that could have pandemic potential. Seasonal influenza, RSV, and COVID-19 may co-circulate throughout the respiratory season and co-infection with multiple viruses at the same time can occur.

Testing

Diagnostic tests available for detection of viruses in respiratory specimens include molecular assays (including rapid molecular assays, reverse transcription polymerase chain reaction (RT-PCR) and other nucleic acid amplification tests); and antigen detection tests (including rapid influenza diagnostic tests and immunofluorescence assays). Sensitivity and specificity can vary by the pathogen or test type, illness onset to specimen collection, the prevalence of viruses in patient population and other factors. Overall, molecular assays have a higher sensitivity and specificity than rapid antigen tests.

When available, multiplex assays for simultaneous detection of influenza, RSV and SARS-CoV-2 viruses should be used. It is possible for a patient to be infected with two or more viruses at the same time. Co-infections can impact the clinical management of acute respiratory illness. Testing for suspected pathogens should be considered particularly in hospitalized patients with severe respiratory disease. Additional guidance for clinicians when SARS-CoV-2 and influenza viruses are co-circulating can be found [here](#).

A negative rapid antigen test does NOT rule out infection and should not be used for treatment or infection control decisions during periods when influenza, RSV, and/or SARS-CoV-2 viruses are known to be circulating.

- RSV: Antigen testing is sensitive in children but less sensitive in adults. Healthcare providers should use highly sensitive rRT-PCR assays when testing older children and adults for RSV.
- COVID-19: A negative viral test result does not rule out infection and should be repeated following CDC and FDA recommendations.
- Influenza: Providers should be aware of circulating influenza viruses. More information is available on the [NC Respiratory Virus Surveillance Dashboard](#).

NC DEPARTMENT OF HEALTH AND HUMAN SERVICES • DIVISION OF PUBLIC HEALTH

LOCATION: 225 North McDowell St., Raleigh, NC 27603
MAILING ADDRESS: 1902 Mail Service Center, Raleigh, NC 27699-1902
www.ncdhhs.gov • TEL: 919-733-7301 • FAX: 919-733-1020

AN EQUAL OPPORTUNITY / AFFIRMATIVE ACTION EMPLOYER

Influenza and COVID-19 testing is available at the North Carolina State Laboratory of Public Health (SLPH). Information on how to submit to SLPH can be found in [SCOPE](#). Specimens should be submitted to SLPH for further testing and characterization in the following circumstances:

1. Specimens from confirmed influenza cases with severe illness and a poor prognosis.
2. Specimens from influenza associated deaths (adult and pediatric).
3. Patients who die with influenza-like illness but have no laboratory evidence of influenza, SARS-CoV-2, or other respiratory infection on a multiplex panel.
4. Patients who are critically ill with influenza-like illness but have no laboratory evidence of influenza, SARS-CoV-2, or other respiratory infection on a multiplex panel.
5. Patients with influenza-like illness, with or without confirmatory testing for influenza, who have had contact with livestock including domestic or wild swine (pigs) or poultry (birds), cattle, sick or dead wild animals, or raw milk or dairy products within the 10 days before symptom onset.
6. Influenza positive specimens that are unable to be subtyped by tests designed to provide an influenza subtyping result.
7. A sample of patients with influenza-like illness seen at facilities participating in the outpatient Influenza-Like Illness Network (ILINet) or Influenza Hospitalization Surveillance Program (IHSP/RESP-NET). [Please consider joining ILINet if you have not done so.](#)

Testing at the SLPH should also be considered for other patients in outbreaks in institutional settings or congregate living facilities and clusters of severe or unusual respiratory illness. Consult [CDC testing and management considerations](#) for residents of long-term care facilities with acute respiratory illness symptoms when SARS-CoV-2 and Influenza viruses are co-circulating as needed.

All specimens submitted to SLPH for influenza or SARS-CoV-2 testing will be tested for both influenza and SARS-CoV-2. Specific guidance regarding specimen collection and transport is available [here](#).

Human Novel or Variant Influenza

NC DPH requests your assistance with enhanced influenza surveillance and testing of individuals presenting with compatible illness and relevant exposure including contact with livestock, poultry, sick or dead wild animals, or raw milk or dairy products within the 10 days before symptom onset. Please follow the algorithm on page 3 for detailed considerations when assessing influenza-like illness in patients. Additional information is available on the [NC DHHS Avian Flu Website](#) including the [Provider Memo page](#).

Availability of Free At-home COVID Test Kits

Community-based organizations and other partners, who are making at-home COVID-19 test kits available to members of their community, can order at-home COVID-19 test kits directly from the federal government for free. Bulk orders can be placed via the Health Partner Order Portal (HPoP). Step-by-step guidance on how to register for an HPoP account and place orders, can be found [here](#). Organizations must not sell or seek reimbursement for the test kits that the federal government provides at no cost.

All U.S. households will be eligible to order four free at-home COVID-19 tests at [COVIDTests.gov](#) at the end of September 2024. Patients can also find organizations providing free at-home COVID-19 test kits

in their community [here](#). There are currently more than 260 active Community Access Points (CAPs), providing free at-home test kits across the state.

Clinicians should contact their [Local Health Departments](#) or the Communicable Disease Branch epidemiologist on call available 24/7 at (919) 733-3419 for questions about respiratory virus testing.

cc: Dr. Erica Wilson, Medical Director, Medical Consultation Unit
Evelyn Foust, Branch Head, Communicable Disease Branch
Dr. Zack Moore, State Epidemiologist

