

NC DEPARTMENT OF HEALTH AND HUMAN SERVICES

ROY COOPER • Governor

MANDY COHEN, MD, MPH • Secretary

MARK T. BENTON • Assistant Secretary for Public Health

Division of Public Health

Developed by the North Carolina Division of Public Health, Communicable Disease Branch

Spotted Fever Group Rickettsiosis Surveillance Summary from 2014—2019

Background

Spotted fever group rickettsioses (SFGR), including Rocky Mountain spotted fever, are a group of bacterial infections caused by *Rickettsia* spp. including *R. rickettsia* and *R. parkeri*. Spotted fevers are transmitted to humans through the bite of an infected tick. In North Carolina the most common vectors of spotted fevers include the American dog tick, *Dermacentor variabilis*, the Rocky Mountain wood tick, *D. andersoni*, and the Lone star tick, *Amblyomma americanum*. The brown dog tick, *Rhipicephalus sanguineus* has been implicated in transmission in other parts of the US. If left untreated, illness can become serious, even leading to death.

Symptomology

Early signs of SFGR are non-specific, including fever and headache. Symptoms may appear 3–12 days following a tick bite. Other signs and symptoms can include nausea, vomiting, stomach pain, muscle pain, lack of appetite, and rash. Rash is a common sign among those infected with *R. rickettsii*, the causative agent of Rocky Mountain spotted fever (RMSF), and usually develops 2-4 days following fever onset. Rashes can look like red splotches or pinpoint dots.

Epidemiology

National

Incidence varies considerably by geographic area. From 2008-2012, 63% of reported SFGR cases originated from five states: Arkansas, Missouri, North Carolina, Oklahoma, and Tennessee¹. Thousands of cases of SFGR occur every year, but it is unknown how many cases are RMSF. Case fatality rates vary annually, but have decreased overall from 28% in 1944 to < 1% in 2001. The national average incidence of SFGR in 2015 was 1.31 cases per 100,000.¹

North Carolina

The number of confirmed and probable cases of spotted fever rickettsiosis has remained stable in North Carolina over the past five years. The highest incidence of SFGR is clustered around central and eastern North Carolina. The 5-year average incidence rate of SFGR in North Carolina from 2014-2018 was 4.85 confirmed and probable cases per 100,000 residents, which is higher than the national average². The incidence of SFGR in North Carolina in 2019 was 6.59 cases per 100,000 residents (using 2018 population data).

Diagnosis

Delay in diagnosis and treatment is the most important factor associated with poor outcomes, and early treatment based on clinical impression is the best way to prevent RMSF progression. Both acute and convalescent serum specimens are needed to confirm the rickettsial infection. Serological tests are often negative during the acute phase of illness, however, physicians may diagnose patients based on the symptoms outlined above. In January 2020, the case definition of SFGR was amended to require an elevated IgG antibody titer of >1:128 within 60 days of illness onset³.

Prevention

Reducing exposure to ticks is the best defense against SFGR. There are a number of methods that can be used to prevent tickborne illness:

- Wear permethrin treated clothing (0.5%) when exploring the outdoors.
- Use EPA registered insect repellents containing DEET or picaridin to deter ticks.
- Avoid ticks in wooded/brushy areas with high grasses and leaf litter by walking in the center of trails.
- Check clothing and skin for ticks you may have encountered while outdoors; shower soon after returning indoors.


Case Demographics

Gender	5 Year Avg (2014-18)		2019	
	No. of Cases	% of total	No. of Cases	% of total
Male	342	70%	453	66%
Female	150	30%	232	34%


Race	5 Year Avg (2014-18)		2019	
	No. of Cases	% of total	No. of Cases	% of total
White	271	56%	410	62%
Black or African American	32	7%	38	6%
Native Hawaiian or Pacific Islander	1	< 1%	0	0%
Amer. Indian or Alaskan	1	< 1%	0	0%
Asian	1	< 1%	6	< 1%
Other	4	< 1%	19	3%
Unknown	179	37%	192	29%

Hispanic Ethnicity	5 Year Avg (2014-18)		2019	
	No. of Cases	% of total	No. of Cases	% of total
Yes	12	3%	28	4%
No	252	55%	380	57%
Unknown	195	42%	257	39%

Confirmed and Probable Spotted Fever Group Rickettsiosis by Year, NC, 2014-2019; n= 3,150




Confirmed and Probable Spotted Fever Group Rickettsiosis by Age Range, NC




Geographic Distribution

Confirmed and Probable Incidence of Spotted Fever Group Rickettsiosis Cases by County of Residence, NC, 2019



Cases by Age

Confirmed and Probable Spotted Fever Group Rickettsiosis cases by Month of Illness Onset, NC



¹Data are based on a national surveillance data found at: https://www.cdc.gov/mmwr/volumes/64/wr/mm6453a1.htm?s_cid=mm6453a1_w

²View NC Disease Statistics here: <https://public.tableau.com/profile/nc.cdb#!/vizhome/NCD3NorthCarolinaDiseaseDataDashboard/DiseaseMapsandTrends>

³CDC Spotted Fever Group Rickettsiosis Case Definition: <https://www.cdc.gov/nndss/conditions/spotted-fever-rickettsiosis/case-definition/2020/>